



Draft Genome Sequence of *Hymenobacter* sp. Strain AT01-02, Isolated from a Surface Soil Sample in the Atacama Desert, Chile

Hansen, Anders Cai Holm; Paulino-Lima, Ivan Glaucio; Fujishima, Kosuke; Rothschild, Lynn Justine; Jensen, Peter Ruhdal

Published in:
Genome Announcements

Link to article, DOI:
[10.1128/genomeA.01701-15](https://doi.org/10.1128/genomeA.01701-15)

Publication date:
2016

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Hansen, A. C. H., Paulino-Lima, I. G., Fujishima, K., Rothschild, L. J., & Jensen, P. R. (2016). Draft Genome Sequence of *Hymenobacter* sp. Strain AT01-02, Isolated from a Surface Soil Sample in the Atacama Desert, Chile. *Genome Announcements*, 4(1), [e01701-15]. <https://doi.org/10.1128/genomeA.01701-15>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Draft Genome Sequence of *Hymenobacter* sp. Strain AT01-02, Isolated from a Surface Soil Sample in the Atacama Desert, Chile

Anders Cai Holm Hansen,^a Ivan Glauco Paulino-Lima,^b Kosuke Fujishima,^c Lynn Justine Rothschild,^d Peter Ruhdal Jensen^a

Microbial Biotechnology & Biorefining, National Food Institute, Technical University of Denmark, Lyngby, Denmark^a; NASA Postdoctoral Program Fellow, NASA Ames Research Center, Moffett Field, California, USA^b; University Affiliated Research Center, NASA Ames Research Center, Moffett Field, California, USA^c; NASA Ames Research Center, Moffett Field, California, USA^d

Here, we report the 5.09-Mb draft genome sequence of *Hymenobacter* sp. strain AT01-02, which was isolated from a surface soil sample in the Atacama Desert, Chile. The isolate is extremely resistant to UV-C radiation and is able to accumulate high intracellular levels of Mn/Fe.

Received 11 December 2015 Accepted 21 December 2015 Published 11 February 2016

Citation Holm Hansen AC, Paulino-Lima IG, Fujishima K, Rothschild LJ, Jensen PR. 2016. Draft genome sequence of *Hymenobacter* sp. strain AT01-02, isolated from a surface soil sample in the Atacama Desert, Chile. *Genome Announc* 4(1):e01701-15. doi:10.1128/genomeA.01701-15.

Copyright © 2016 Holm Hansen et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Anders Cai Holm Hansen, achh@food.dtu.dk.

Hymenobacter sp. strain AT01-02, isolated from the Atacama Desert, Chile, is a Gram-negative pink-pigmented bacterium that thrives in an extremely dry environment, where it is exposed to large temperature variations and high levels of solar UV radiation (1, 2). Several *Hymenobacter* spp. were reported to be radiation resistant (3–6), and isolate AT01-02 is more resistant than *Deinococcus radiodurans* to UV-C irradiation, being able to accumulate high intracellular levels of Mn/Fe (our unpublished data).

Here, we describe a draft genome sequence of *Hymenobacter* sp. AT01-02 to investigate the genetic mechanisms involved in the survival of this organism in such an extreme environment. Total genomic DNA was extracted from colonies using the Microbial DNA isolation kit (Mo Bio Laboratories, Solano Beach, CA, USA), according to the manufacturer's instructions. The genome was sequenced on an Ion Torrent instrument, producing 2,680,302 reads, which were trimmed with respect to quality and size (cutoff, Q20; size, 50 to 270 bp). The trimmed reads were *de novo* assembled using CLC bio Genomics Workbench into 272 contigs, constituting a total genome size of 5,043,991 bp, with sizes ranging from 517 bp to 315,336 bp. The average size was 18,544 bp, and the N_{50} was 74,558 bp. The G+C content was 55.5%, with a genome coverage of 90%.

We annotated the assembled genome using the IMG-ER portal (<https://img.jgi.doe.gov/cgi-bin/er/main.cgi>). We found 50 predicted RNAs, including 4 rRNAs, 42 tRNAs, and 4 miscellaneous RNAs. The RAST result showed 4,821 protein-coding genes (coding sequences), including 1,360 known and 3,461 unknown subsystems, with 759 protein-coding genes connected to KEGG pathways.

We found 35 genes related to stress response, including 24 genes for oxidative stress, 5 genes for cold shock, 5 genes for general stress, and 1 gene for osmotic stress. We also found 21 genes related to the biosynthesis of secondary metabolites, 14 genes related to drug resistance, and 1 multidrug resistance gene (efflux pump membrane protein). Additionally, we identified 38 genes for the metabolism of terpenoids and polyketides, 29 genes for

xenobiotic biodegradation and metabolism, 43 genes related to DNA replication and repair (UvrABC system, *recA*, and *MutS*), and 29 genes related to membrane transport (including 16 ABC transporters, 11 secretory systems, and 2 manganese transporters of the natural resistance-associated macrophage protein [NRAMP] family). Similar to what was reported recently for a different *Hymenobacter* isolate (7), we found 4 teichoic and 3 lipoteichoic biosynthesis genes, which are characteristically present in Gram-positive organisms, although AT01-02 is Gram negative. Interestingly, 16 genes indicated the presence of bacteriophage (including 9 integrase, 5 transposase, 1 terminase, and 1 lamin tail domain), which is related to low optical densities obtained with liquid cultures, and to a clear center observed in old colonies grown on agar plates. Also, 2 genes were related to exopolysaccharide biosynthesis, which is connected to morphological aspects of the colonies grown on plates, including brilliance and stickiness.

The genome of AT01-02 exhibits a diverse suite of stress-responsive and pigment-producing genes, along with bacteriophage genes and many genes for the biosynthesis of secondary metabolites and biodegradation of xenobiotics. These characteristics represent its potential for industrial applications and will provide insights to better understand the survival mechanisms against environmental stresses, such as desiccation and UV radiation.

Nucleotide sequence accession numbers. This whole-genome shotgun project has been deposited at DDBJ/EMBL/GenBank under the accession number [JZIR000000000](https://www.ncbi.nlm.nih.gov/nuclot/JZIR000000000). The version described in this paper is version JZIR020000000.

ACKNOWLEDGMENTS

We thank the DMAC facility (DTU Microarray Core, Technical University of Denmark, Lyngby, Denmark) for assisting with sample preparation and providing sequencing data. We also thank Armando Azua-Bustos for sampling at the Atacama Desert.

This work was supported by I.G.P.-L.'s appointment to the NASA

Postdoctoral Program at NASA Ames Research Center, administered by Oak Ridge Associated Universities through a contract with NASA.

REFERENCES

1. Azua-Bustos A, Caro-Lara L, Vicuña R. 2015. Discovery and microbial content of the driest site of the hyperarid Atacama Desert, Chile. *Environ Microbiol Rep* 7:388–394.
2. Paulino-Lima IG, Azua-Bustos A, Vicuña R, González-Silva C, Salas L, Teixeira L, Rosado A, da Costa Leitao AA, Lage C. 2013. Isolation of UVC-tolerant bacteria from the hyperarid Atacama Desert, Chile. *Microb Ecol* 65:325–335. <http://dx.doi.org/10.1007/s00248-012-0121-z>.
3. Collins MD, Hutson RA, Grant IR, Patterson MF. 2000. Phylogenetic characterization of a novel radiation-resistant bacterium from irradiated pork: description of *Hymenobacter actinosclerus* sp. nov. *Int J Syst Evol Microbiol* 50:731–734. <http://dx.doi.org/10.1099/00207713-50-2-731>.
4. Dai J, Wang Y, Zhang L, Tang Y, Luo X, An H, Fang C. 2009. *Hymenobacter tibetensis* sp. nov., a UV-resistant bacterium isolated from Qinghai-Tibet plateau. *Syst Appl Microbiol* 32:543–548. <http://dx.doi.org/10.1016/j.syapm.2009.09.001>.
5. Lee J-J, Srinivasan S, Lim S, Joe M, Lee SH, Kwon SA, Kwon YJ, Lee J, Choi JJ, Lee HM, Auh YK, Kim MK. 2014. *Hymenobacter swuensis* sp. nov., a gamma-radiation-resistant bacteria isolated from mountain soil. *Curr Microbiol* 68:305–310. <http://dx.doi.org/10.1007/s00284-013-0478-3>.
6. Zhang Q, Liu C, Tang Y, Zhou G, Shen P, Fang C, Yokota A. 2007. *Hymenobacter xinjiangensis* sp. nov., a radiation-resistant bacterium isolated from the desert of Xinjiang, China. *Int J Syst Evol Microbiol* 57:1752–1756. <http://dx.doi.org/10.1099/ijs.0.65033-0>.
7. Koo H, Ptacek T, Crowley M, Swain AK, Osborne JD, Bej AK, Andersen DT. 2014. Draft genome sequence of *Hymenobacter* sp. strain IS2118, isolated from a freshwater lake in Schirmacher Oasis, Antarctica, reveals diverse genes for adaptation to cold ecosystems. *Genome Announc* 2(4): e00739-14. <http://dx.doi.org/10.1128/genomeA.00739-14>.